




INVESTOR IN PEOPLE

The Patent Office
 Concept House
 Cardiff Road
 Newport
 South Wales
 NP10 8QQ



PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
 COMPLIANCE WITH RULE 17.1(a) OR (b)

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed *Stephen Hordley*
 Dated 23 July 2004

Best Available Copy



Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office
Cardiff Road
Newport
Gwent NP9 1RH

1. Your reference

LVK/HG/PB60434P

2. Patent application number

(The Patent Office will fill in his part)

0318728.3

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Glaxo Group Limited
Glaxo Wellcome House, Berkeley Avenue,
Greenford, Middlesex UB6 0NN, Great Britain

Patents ADP number (if you know it)

473587003

If the applicant is a corporate body, give the country/state of its incorporation

UK

United Kingdom

4. Title of the invention

Process

5. Name of your agent (if you have one)

Corporate Intellectual Property

"Address for service" in the United Kingdom to which all correspondence should be sent
(including the postcode)

GlaxoSmithKline
Corporate Intellectual Property (CN9 25.1)
980 Great West Road
BRENTFORD
Middlesex TW8 9GS

Patents ADP number (if you know it)

7960982003

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or each of these earlier applications and (if you know it) the or each application number

Country Priority application number Date of filing
(if you know it) (day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer yes if:

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is named as an applicant, or
- c) any named applicant is a corporate body

See note (d)

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form.
Do not count copies of the same document

Continuation sheets of this form	13
Description	1
Claim(s)	
Abstract	
Drawings	

10. If you are also filing any of the following, state how many against each item.

Priority Documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patents Form 9/77*)

Request for substantive examination (*Patents Form 10/77*)

Any other documents
(please specify)

11.

We request the grant of a patent on the basis of this application

Signature

L V Knight

Date 7-Aug-03

12. Name and daytime telephone number of person to contact in the United Kingdom

L V Knight 01279 644644

Warning

After an application for a Patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission unless an application has been filed at least six weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505*
- Write your answers in capital letters using black ink or you may type them.*
- If there is not enough space for all relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.*
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.*
- For details of the fee and ways to pay please contact the Patent Office.*

Process

The present invention relates to a process for preparing radiolabelled compounds. More specifically, the present invention relates to a process for preparing radiolabelled compounds, which involves incorporation of radioactive carbonyl groups into precursors, which are then used to make the radiolabelled compounds. These radiolabelled compounds have a number of uses including in vivo imaging techniques such as positron emission tomography.

Positron emission tomography (PET) is a non-invasive imaging technique that offers high spatial and temporal resolution and allows quantification of tracer concentrations in tissues. The technique involves the use of radiotracers labelled with positron emitting radionuclides, which permit measurement of parameters regarding the physiology or biochemistry of a variety of living tissues.

Compounds can be labelled with positron or gamma emitting radionuclides. The most commonly used positron emitting (PET) radionuclides are ^{11}C , ^{18}F , ^{15}O and ^{13}N , which are accelerator produced, and have half lives of 20.4, 109.8, 2 and 10 minutes respectively. Due to their short half-lives ^{11}C , ^{15}O and ^{13}N labelled radiopharmaceuticals have to be used at the site of production and require the development of specific synthetic procedures.

^{11}C ($T_{1/2}=20.4\text{min}$) is an important neutron-deficient radionuclide for PET because it can substitute for non-radioactive carbon in any organic molecule without altering their biological and physiochemical properties. An important part of the elaboration of new procedures to incorporate PET radionuclides into molecules is the development and handling of new ^{11}C labelled precursors.

^{11}C can be produced in the absence of the naturally occurring stable isotopes ^{12}C and ^{13}C , and with high yields on a small proton accelerator using the $^{14}\text{N}(p,\alpha)^{11}\text{C}$ reaction in a target gas containing nitrogen (Christman, et al., 1975; Clark, et al., 1975 and Welch et al., 1968). In the presence of oxygen trace (e.g. 0.1% oxygen), the radiochemical species formed is $[^{11}\text{C}]$ carbon dioxide which is suitable for use directly as in the ^{11}C -carboxylation of Grignard reagents (organomagnesium halides). $[^{11}\text{C}]$ carbon dioxide can also be converted into a variety of secondary radiolabelled chemical entities such as high specific activity $[^{11}\text{C}]$ methyl iodide.

An important consideration for radiolabelling with carbon-11 is the maximization of specific activity of the radiolabelled compound. Isotopic dilution of $[^{11}\text{C}]$ carbon dioxide with atmospheric carbon dioxide (3.4×10^4 ppm) substantially reduces its specific activity and therefore limits the application of the resultant radiolabelled compound as a PET probe.

As an alternative method to using $[^{11}\text{C}]$ carbon dioxide for radiolabelling compounds, $[^{11}\text{C}]$ carbon monoxide may be used instead, as it is less prone to isotopic dilution with

atmospheric carbon monoxide (0.1 ppm). Methods for the production of [^{11}C]carbon monoxide by reducing [^{11}C]carbon dioxide using reducing metals at high temperatures are well known (Gmelins 1972; Clark, et al., 1975; Zeisler, et al., 1997). Zinc and molybdenum are the most widely used reducing agent for the [^{11}C]carbon dioxide/carbon monoxide conversion.

However, it is difficult to trap ^{11}CO in the small volume of organic solvent in which most of the precursors for the production of radiolabelled compounds are soluble. Small volumes of solvent are required because this allows easy isolation of the radiolabelled product by means of preparative HPLC and increases the concentration of the starting materials in the reaction mixture, thereby forcing the reaction in the desired direction.

In 1978 Roeda, et al., described a method for the production of [^{11}C]phosgene from [^{11}C]carbon monoxide however, its practical use in the production of radiopharmaceuticals has been very limited due low yields and the lack of suitable equipment and methods to efficiently trap and react carbon monoxide.

Existing methods for the trapping of [^{11}C]carbon monoxide for the production of radiolabelled compounds rely on the use of high pressure or recirculation of [^{11}C]carbon monoxide to maintain adequately high levels of [^{11}C]carbon monoxide in solution (Kihlberg, et al., 1999; Hostetler, et al., 2002). This requires the use of dedicated automated robotic systems for the handling of [^{11}C]carbon monoxide and specialised equipment.

Borane carbonyl (H_3BCO) is the immediate precursor to boranocarbonates, such as the potassium salt $\text{K}_2[\text{H}_3\text{BCO}_2]$ which were reported to release CO in water at elevated temperatures in 1967 (Malone et al., 1967; Malone et al., 1967a). Although yields of the solid, air stable $\text{K}_2[\text{H}_3\text{BCO}_2]$, produced from the known methods of $\text{B}_2\text{H}_6 + \text{CO}$ are good, it is not convenient to work under pressurised conditions with H_3BCO , as it is a pyrophoric gas (Carter, et al., 1965; Mayer, 1971). Alberto et al., (2001) found that by preparing H_3BCO from commercially available $\text{H}_3\text{B.THF}$ and reacting it in situ with an alcoholic solution of potassium hydroxide, $\text{K}_2[\text{H}_3\text{BCO}_2]$ could be produced at ambient pressures. This result was achieved by controlling the equilibrium of the two-way reaction between H_3BCO and $\text{H}_3\text{B.THF}$ by selectively condensing the THF out of the reaction. The resultant $\text{K}_2[\text{H}_3\text{BCO}_2]$ was then used as an in situ source of CO in aqueous solution and as a reducing agent.

It has now been found that radiolabelled H_3BCO can be used to release radiolabelled carbon monoxide in organic solvents, aqueous solvents and mixtures of organic and aqueous solvents in order to prepare radiolabelled compounds without the need for high pressure autoclaves or recirculation units.

Accordingly, in a first aspect the invention provides a process for the preparation of radiolabelled H_3BCO comprising contacting H_3B in a suitable solvent with carbon monoxide, characterised in that the carbon monoxide is radiolabelled.

- 5 Radiolabelled H_3BCO may be prepared by the reaction of borane (H_3B) in a suitable solvent with radiolabelled carbon monoxide. Suitable solvents for this reaction are those which solubilize H_3B and allow it to co-ordinate with free electron pairs of the oxygen, for example tetrahydrofuran (THF) and ethers such as diethyl ether and dioxane. THF is preferred as a solvent due to its physical characteristics of a high boiling point, a lower
10 affinity towards water and its comparable low price.

- Hydrides of other elements, such as aluminium gallium, indium and thallium hydride would also be expected to co-ordinate with radiolabelled carbon monoxide. However, the instability of aluminium hydride in solvents suitable for this reaction means that if an
15 aluminium compound were to be used it would preferably be compounds such as AlCl_3 in THF or aluminium tri organyls.

- Free solvent may be removed from the reaction by condensation or other suitable means such as a solid support. This achieves the advantage of shifting the equilibrium of the
20 reaction towards increased production of radiolabelled H_3BCO .

The carbon monoxide used in the reaction may be labelled by any conventional method with any of the following isotopes ^{11}C , ^{13}C , ^{14}C or ^{18}O . Preferably ^{11}C is used.

- 25 Suitable solvents for use in the process of the invention include ethers such as diethyl ether and dioxane, and tetrahydrofuran. Preferably tetrahydrofuran is used.

- In a second aspect the invention provides the use of radiolabelled H_3BCO prepared according to the first aspect of the invention, as a donor of radiolabelled carbon monoxide
30 in the manufacture by carbonylation of radiolabelled compounds.

- In practice the second aspect of the invention may be carried out by using the radiolabelled H_3BCO prepared according to the first aspect of the invention in a coupling
35 reaction as set out in Scheme 1 below, in which coupling reactions are typically carried out with a halide or a triflate (trifluoromethanesulfonate) with a nucleophile (alcohol, amine, thiol) or a organostannane, a base and a catalyst such as a palladium(0) catalyst to obtain esters, amides, ketones, aldehydes, carboxylic thioesters or by reacting a nitro
40 component or an azido derivative to form isocyanate derivatives or condensing two nucleophiles in presence of selenium to synthesized carbamates, thiocarbamates, carbonates and ureas.


The starting materials and reagents for use in the first and second aspects of the invention are available commercially or can be synthesised by well-known and

conventional methods. The reaction conditions used in the formation of non-radiolabelled H_3BCO can be sourced from Alberto et al., (2001), other reaction conditions such as the radiolabelling of CO and carbonylation reactions are well known.

- 5 $[^{11}\text{C}]\text{CO}$, prepared by reduction of $[^{11}\text{C}]\text{CO}_2$ with a reducing metal (commonly zinc or molybdenum), is trapped using conventional methods such as molecular sieves in liquid nitrogen or silica and is then carried into a solution of $\text{BH}_3\cdot\text{THF}$ using an inert gas carrier. The $[^{11}\text{C}]\text{borane carbonyl}$ ($[^{11}\text{C}][\text{H}_3\text{BCO}]$) complex thus formed is then carried through to a reaction chamber in which it is reacted with suitable components to construct the required compound using conventional coupling methods. Conventional coupling reaction often
10 take place at elevated temperatures and the reaction chamber may be made of materials suitable for use in a microwave (such as glass).




- 15 In order to promote the formation of the $[^{11}\text{C}]\text{borane carbonyl}$ THF is removed from the reaction, typically by condensation. Coupling reactions are typically carried out reacting $[^{11}\text{C}]\text{borane carbonyl}$ with the appropriate starting materials and reagents as depicted in scheme 1.

(I) Formation of radioactive BH_3CO




(II) Coupling reaction

(a) when R1 and R2 are aryl, vinyl or alkyl;
 X is I, Br, Cl or OTf; and
 Y is NH_2 , NHR , OH , SH or
 $\text{Sn}(\text{R}_3)_3$ (with R_3 = alkyl or aryl)




or (b) when X and Y are O, N or S
 and R1 and R2 are aryl or alkyl



Scheme1

- 5 Suitable compounds for radiolabelling by this method are those which contain a carbonyl group (some examples are shown in Scheme 2).



Scheme 2

5 Amides and imides can also contain lactams and carboxylic esters can also contain lactones.

In a third aspect the invention provides radiolabelled H_3BCO prepared in accordance with the first aspect of the invention.

10

In fourth aspect the invention provides radiolabelled compounds prepared by carbonylation in accordance with the second aspect of the invention.

15

Edidepride (*N*-((*S*)-1-Ethyl-pyrrolidin-2-ylmethyl)-3-iodo-5-methoxy-benzamide), FLB (5-bromo-*N*-((*S*)-1-ethyl-pyrrolidin-2-ylmethyl)-2,3-dimethoxy-benzamide) and raclopride (3,5-dichloro-*N*-((*S*)-1-ethyl-pyrrolidin-2-ylmethyl)-2-hydroxy-6-methoxy-benzamide), which are all dopamine D2 ligands and PK11195 (1-(2-Chloro-phenyl)-isoquinoline-3-carboxylic acid), which is a benzodiazepine receptor ligand are commonly used PET ligands that contain carbonyl groups that can be labelled with $^{11}\text{C}[\text{CO}]$.

20

In a fifth aspect the invention provides use of the radiolabelled compounds according to the fourth aspect of the invention in imaging techniques such as positron emission tomography, modified single photon emission tomography and autoradiography (classical and phosphor imaging plates).

25

In a sixth aspect the invention provides a composition comprising a radiolabelled compound in accordance with the fourth aspect of the invention and a pharmaceutically acceptable carrier or carriers, suitable for use in the above mentioned imaging techniques.

30

Examples

Abbreviation list:


THF: Tetrahydrofuran

TEA: Triethylamine

DBU: 1,8-Diazabicyclo[5.4.0]undec-7-ene

TMP: Tetramethylpiperidine

DMF: Dimethylformamide

10 Synthesis of [^{11}C]N-benzyl-benzamide

Example 1

15 Preparation of the reaction vial

Palladium(II) diacetate (0.5 mg, 0.0022 mmol) and triphenylphosphine (2.9 mg, 0.011 mmol) were dissolved in 700 μL THF (degassed by bubbling N_2 through it for few minutes). Then, iodobenzene (1.5 mg, 0.00735 mmol) and benzylamine (1.2 mg, 0.011 mmol) dissolved in 300 μL THF (degassed by bubbling N_2 through it for few minutes) were added to the solution of palladium complex. TEA (1.6 μL , 0.0088 mmol) was added, and the reaction vial was placed in the reaction-setup in a bath at -78°C .

Synthesis

[^{11}C]Carbon dioxide was produced by the $^{14}\text{N}(\text{p},\alpha)^{11}\text{C}$ nuclear reaction using a nitrogen gas target (containing 1% oxygen) pressurised to 150 psi and bombarded with 16 MeV protons using the General Electric Medical Systems PETtrace 200 cyclotron. Typically, the irradiation time was 30 minutes using a 40 μA beam current. After irradiation, [^{11}C]carbon dioxide was trapped and concentrated on 4Å molecular sieves. The trapped [^{11}C]CO₂ was released from molecular sieves in a stream of nitrogen (30 mL/min) by heating them to 350°C . [^{11}C]CO₂ was reduced on-line to [^{11}C]carbon monoxide after passing through a quartz tube filled with zinc granular heated to 400°C . The produced [^{11}C]carbon monoxide was transferred in our system set-up at 30 mL/min, where it was condensed on 4Å molecular sieves at -196°C . After 6 min delivery and trapping of the [^{11}C]CO, the radioactive gas was then released at room temperature in a flow of nitrogen (6 mL/min) to bubble through a $\text{BH}_3\cdot\text{THF}$ solution (1.5 mL of a 1.0 M solution) in order to make the [^{11}C]BH₃·CO complex. This complex was carried with the flow of nitrogen through an empty vial cooled at -60°C to remove the THF, and finally through the reaction vial containing the reactants (cf. preparation of the reaction vial above) cooled at -78°C .

The trapping process took approximately 6 mins (when the radioactivity level measured in the reaction vial has reached a maximum). The delivery tubings were then removed and the reaction vial heated in an oven at 110°C for 10 mins. The crude product was filtered through a 0.45 µm filter and analysed using analytical radio HPLC.

Analytical HPLC was performed using a Dionex system (SUMMIT HPLC system), equipped with a Dionex HPLC pump (Model P 680A LPG) with a 200 µl injection loop connected in series with a Phenomenex Spherclone 5u ODS(2) column (250 x 4.60 mm, 5 µm), a variable Dionex UV/VIS detector (Type UVD 170U/340U) in series with a sodium iodide radiodetector of in-house design.

The desired end-product was identified by co-injection with a non-radioactive reference. The given yields of the product are based on the final radioactivity trapped in the reaction vial at EOS (End Of Synthesis).

The analytical HPLC showed the formation of the desired radiolabelled [¹¹C]*N*-benzylbenzamide in Example 1 in approximately 1.7% yield.

Example 2

The synthesis of [¹¹C]*N*-benzyl-benzamide was carried out as described in Example 1 except that the palladium(II) diacetate (0.5 mg, 0.0022 mmol) and triphenylphosphine (2.9 mg, 0.011 mmol) were dissolved in 700 µL of a solution of THF:H₂O, 4:1 (degassed by bubbling N₂ through it for few minutes), the iodobenzene (1.5 mg, 0.00735 mmol) and benzylamine (1.2 mg, 0.011 mmol) were dissolved in 300 µL of a solution of THF:H₂O, 4:1 (degassed by bubbling N₂ through it for few minutes). The reaction vial was placed in the reaction-setup in a bath at 0°C and after the trapping of the [¹¹C]BH₃.CO the reaction vial was heated at 120°C for 8 mins. The analytical HPLC showed the formation of the desired radiolabelled [¹¹C]*N*-benzylbenzamide in approximately 7% yield.

Example 3

The synthesis of [¹¹C]*N*-benzyl-benzamide was carried out as described in Example 1 except that the palladium(II) diacetate (0.5 mg, 0.0022 mmol) and triphenylphosphine (2.9 mg, 0.011 mmol) were dissolved in 700 µL of a solution of THF + 2% H₂O (degassed by bubbling N₂ through it for few minutes), the iodobenzene (1.5 mg, 0.00735 mmol) and benzylamine (1.2 mg, 0.011 mmol) were dissolved in 300 µL of a solution of THF + 2% H₂O (degassed by bubbling N₂ through it for few minutes) and after the trapping of the [¹¹C]BH₃.CO, the reaction vial was heated at 120°C for 8 mins. The analytical HPLC showed the formation of the desired [¹¹C]*N*-benzylbenzamide in approximately 30% yield.

Example 4

The synthesis of [¹¹C]*N*-benzyl-benzamide was carried out as described in Example 1 except that the palladium(II) diacetate (0.5 mg, 0.0022 mmol) and triphenylphosphine (2.9 mg, 0.011 mmol) were dissolved in 700 µL of a solution of THF + 1% H₂O (degassed by

bubbling N₂ through it for few minutes), the iodobenzene (1.5 mg, 0.00735 mmol) and benzylamine (1.2 mg, 0.011 mmol) were dissolved in 300 µL of a solution of THF + 1% H₂O (degassed by bubbling N₂ through it for few minutes) and after the trapping of the [¹¹C]BH₃.CO the reaction vial was heated at 50°C for 8 mins. The analytical HPLC showed the formation of the desired [¹¹C]N-benzylbenzamide in approximately 17% yield.

Example 5

The synthesis of [¹¹C]N-benzyl-benzamide was carried out as described in Example 4 except that after the trapping of the [¹¹C]BH₃.CO, the reaction vial was heated at 70°C for 8 mins. The analytical HPLC showed the formation of the desired [¹¹C]N-benzylbenzamide in approximately 47% yield.

Example 6

The synthesis of [¹¹C]N-benzyl-benzamide was carried out as described in Example 4 except that after the trapping of the [¹¹C]BH₃.CO, the reaction vial was heated at 85°C for 8 mins. The analytical HPLC showed the formation of the desired [¹¹C]N-benzylbenzamide in approximately 47% yield.

Example 7

The synthesis of [¹¹C]N-benzyl-benzamide was carried out as described in Example 4 except that after the trapping of the [¹¹C]BH₃.CO, the reaction vial was heated at 120°C for 8 mins. The analytical HPLC showed the formation of the desired [¹¹C]N-benzylbenzamide in approximately 47% yield.

Example 8

The synthesis of [¹¹C]N-benzyl-benzamide was carried out as described in Example 4 except that after the trapping of the [¹¹C]BH₃.CO, the reaction vial was heated at 140°C for 8 mins. The analytical HPLC showed the formation of the desired [¹¹C]N-benzylbenzamide was approximately 28% yield.

Example 9

The synthesis of [¹¹C]N-benzyl-benzamide was carried out as described in Example 5 except that DBU (1.3 µL, 0.0016 mmol) was used instead of TEA. The analytical HPLC showed traces of the formation of the desired [¹¹C]N-benzylbenzamide.

Example 10

The synthesis of [¹¹C]N-benzyl-benzamide was carried out as described in Example 5 except that 2,2,6,6-TMP (1.7 µL, 0.009 mmol) was used instead of TEA. The analytical HPLC showed the formation of the desired [¹¹C]N-benzylbenzamide in approximately 8% yield.

Example 11

The synthesis of [^{11}C]N-benzyl-benzamide was carried out as described in Example 5 except that pyridine (0.7 μL , 0.0088 mmol) was used instead of triethylamine and the reaction vial was heated from 40 to 80°C for 15 mins. The analytical HPLC showed the formation of the desired [^{11}C]N-benzylbenzamide in approximately 28% yield.

Example 12


The synthesis of [^{11}C]N-benzyl-benzamide was carried out as described in Example 5 except that benzylamine (3.6 mg, 0.034 mmol) was used instead of TEA and the reaction vial was heated 90°C for 8 mins. The analytical HPLC showed the formation of the desired [^{11}C]N-benzylbenzamide in approximately 20% yield.

Example 13

The synthesis of [^{11}C]N-benzyl-benzamide was carried out as described in Example 4 except that the palladium(II) diacetate, triphenylphosphine, iodobenzene and benzylamine benzylamine were dissolved in DMF, and after the addition of TEA the reaction vial was placed in the reaction-setup in a bath at -50°C. After the trapping of the [^{11}C]BH₃.CO the reaction vial was heated at 90°C for 8 mins. The analytical HPLC showed the formation of the desired [^{11}C]N-benzylbenzamide in approximately 23% yield.

Example 14

The synthesis of [^{11}C]N-benzyl-benzamide was carried out as described in Example 4 except that the palladium(II) diacetate, triphenylphosphine, iodobenzene and benzylamine benzylamine were dissolved in 1,2-dichloroethane, and after the addition of TEA the reaction vial was placed in the reaction-setup in a bath at -20°C. After the trapping of the [^{11}C]BH₃.CO the reaction vial was heated at 110°C for 8 mins. The analytical HPLC showed the formation of the desired [^{11}C]N-benzylbenzamide in approximately 12% yield.

Synthesis of [^{11}C]phthalide**Example 15**

Tetrakis(triphenylphosphine)palladium(0) (1.1 mg, 0.95 μmol) was dissolved in 500 μL of a solution of THF + 1% H₂O (degassed by bubbling N₂ through it for few minutes). Then, a mixture of 2-bromobenzyl alcohol (1.1 mg, 0.006 mmol) and K₂CO₃ (5 mg, 0.036 mmol) were dissolved in 300 μL of THF + 1% H₂O (degassed by bubbling N₂ through it for few minutes) and added to the solution of the palladium complex. The reaction vial was placed in the reaction-setup in a bath at -78°C. After the trapping of the [^{11}C]BH₃.CO as described

in Example 1, the reaction was heated at 100°C for 4 mins. The analytical HPLC showed the formation of the desired [¹¹C]phthalide in traces.

Example 16

- 5 Palladium(II) diacetate (0.8 mg, 0.0035 mmol) and triphenylphosphine (5 mg, 0.020 mmol) were dissolved in 700 µL of a solution of THF + 1% H₂O (degassed by bubbling N₂ through it for few minutes). Then, a mixture of 2-bromobenzyl alcohol (2.2 mg, 0.012 mmol) and K₂CO₃ (5 mg, 0.036 mmol) were dissolved in 300 µL of THF + 1% H₂O (degassed by bubbling N₂ through it for few minutes) and added to the solution of the
10 palladium complex. The reaction vial was placed in the reaction-setup in a bath at -78°C. After the trapping of the [¹¹C]BH₃.CO as described in Example 1, the reaction was heated at 120°C for 5 mins. The analytical HPLC showed the formation of the desired [¹¹C]phthalide in traces.

Example 17

- 15 Palladium(II) diacetate (0.8 mg, 0.0035 mmol) and triphenylphosphine (5 mg, 0.020 mmol) were dissolved in 700 µL of a solution of THF (degassed by bubbling N₂ through it for few minutes). Then, a mixture of 2-bromobenzyl alcohol (2.2 mg, 0.012 mmol) and DBU (2.0 µL, 0.014 mmol) was dissolved in 300 µL of THF (degassed by bubbling N₂ through it for
20 few minutes) and added to the solution of the palladium complex. The reaction vial was placed in the reaction-setup in a bath at -78°C. After the trapping of the [¹¹C]BH₃.CO as described in Example 1, the reaction was heated at 110°C for 5 mins. The analytical HPLC showed the formation of the desired [¹¹C]phthalide in traces.

Example 18

- 25 Palladium(II) diacetate (0.8 mg, 0.0035 mmol) and triphenylphosphine (5 mg, 0.020 mmol) were dissolved in 700 µL of a solution of THF (degassed by bubbling N₂ through it for few minutes). Then, a solution of 2-bromobenzyl alcohol (2.2 mg, 0.012 mmol) in 300 µL of THF (degassed by bubbling N₂ through it for few minutes) was added to the solution of the
30 palladium complex. The reaction vial was placed in the reaction-setup in a bath at -78°C. After the trapping of the [¹¹C]BH₃.CO as described in Example 1, the reaction was heated at 120°C for 5 mins. The analytical HPLC showed the formation of the desired [¹¹C]phthalide in approximately 40% yield.

Example 19

- 35 Palladium(II) diacetate (0.5 mg, 0.0022 mmol) and triphenylphosphine (2.9 mg, 0.011 mmol) were dissolved in 700 µL of a solution of THF + 1% H₂O (degassed by bubbling N₂ through it for few minutes). Then, a mixture of 2-bromobenzyl alcohol (2.2 mg, 0.012 mmol) and TEA (1.9 µL, 0.014 mmol) were dissolved in 300 µL of THF + 1% H₂O
40 (degassed by bubbling N₂ through it for few minutes) and added to the solution of the palladium complex. The reaction vial was placed in the reaction-setup in a bath at -78°C. After the trapping of the [¹¹C]BH₃.CO as described in Example 1, the reaction was heated

at 90°C for 8 mins. The analytical HPLC showed the formation of the desired [¹¹C]phthalide in approximately 26% yield.

Example 20

- 5 Palladium(II) diacetate (1.0 mg, 0.0044 mmol) and triphenylphosphine (6 mg, 0.022 mmol) were dissolved in 700 μ L of a solution of THF + 1% H₂O (degassed by bubbling N₂ through it for few minutes). Then, a mixture of 2-bromobenzyl alcohol (2.2 mg, 0.012 mmol) and TEA (1.9 μ L, 0.014 mmol) were dissolved in 300 μ L of THF + 1% H₂O (degassed by bubbling N₂ through it for few minutes) and added to the solution of the
- 10 palladium complex. The reaction vial was placed in the reaction-setup in a bath at -78°C. After the trapping of the [¹¹C]BH₃.CO as described in Example 1, the reaction was heated at 90°C for 8 mins. The analytical HPLC showed the formation of the desired [¹¹C]phthalide in approximately 20% yield.

15

References

Alberto et al *J. Am. Chem. Soc.* (2001) 123, 3135-3136

5 Carter, J. C., Parry R. W., *J. Am. Chem. Soc.* (1965), 87, 2354-2358

Christman D.R., Finn R.D., Karlstrom K. I. and Wolf A. P. (1975) *Int. J. Appl. Radiat. Isot.* 26, 435-442

10 Clark, J.C. and Buckingham, P.D. (1975) *Short-lived Radioactive Gases for Medical Use*, p231. Butterworths, London

Gmelins (1972) *Handbuch der Anorganischen Chemie* Vol. 'Kohlenstoff' C2, p.203. Springer, Heidelberg

15 Hostetler, E.D. and Burns, H.D., *Nucl. Med. Biol.* (2002) 29, 845-848

Kihlberg, T., Bengt Langstrom T.B., *J. Org. Chem.* (1999) 64, 9201-9205

20 Malone L. J., Parry R. W., *Inorg. Chem.* (1967), 6, 817-822

Malone L. J., *Inorg. Chem.* (1967a), 6, 2260-2262

Mayer E., *Monatsh. Chem.* (1971), 102, 940-945

25 Roeda D., Crouzel C. and Van Zanten B (1978) *Radiochem. Radioanal. Letts* 33, 175-178

Welch M.J. and Ter-Pogossian M.M. (1968) *Radiation Res.* 36, 580-587)

30 Zeisler S.K., Nader M., Theobald A. and Oberdorfer F. (1997) *Appl. Radiat. Isot.* vol. 48, 1091-1095

Claims

1. A process for the preparation of radiolabelled H_3BCO comprising contacting H_3B in a suitable solvent with carbon monoxide, characterised in that the carbon monoxide is radiolabelled.
2. A process according to claim 1, wherein production of H_3BCO is promoted by removal of free solvent from the mixture.
3. A process according to claim 2, wherein removal of free solvent from the mixture is achieved by condensation.
4. A process according to any one of claims 1 to 3, wherein a suitable solvent is selected from any ether or tetrahydrofuran.
5. A process according to any one of claims 1 to 4, wherein a suitable solvent is selected from diethyl ether, dioxane or tetrahydrofuran.
6. A process according to any one of claims 1 to 5, wherein a suitable solvent is tetrahydrofuran.
7. A process according to any one of claims 1 to 6, wherein the carbon monoxide is radiolabelled with ^{11}C , ^{13}C , ^{14}C or ^{18}O .
8. A process according to claim 7, wherein the radiolabel is ^{11}C .
9. A process for preparing radiolabelled compounds by carbonylation using radiolabelled H_3BCO prepared according to any one of claims 1 to 8 as a donor of radiolabelled carbon monoxide.
10. A radiolabelled compound prepared using a process according to claim 9.
11. Use of a radiolabelled compound according to claim 10 in imaging techniques.
12. Use according to claim 11, wherein the imaging technique is selected from positron emission tomography, modified single photon emission tomography or autoradiography.
13. Use according to claim 11, wherein the imaging technique is selected from positron emission tomography.
15. A product of a process according to any one of claims 1 to 9.
16. A composition comprising a radiolabelled compound according to claim 10.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.